

1/28





FIGURE 1



Probe	R^1	R^2	R^3	R^4
<i>o</i> -BMOQBA	OCH_3	B(OH)_2	H	H
<i>m</i> -BMOQBA	OCH_3	H	B(OH)_2	H
<i>p</i> -BMOQBA	OCH_3	H	H	B(OH)_2
BMOQ	OCH_3	H	H	H
<i>o</i> -BMQBA	CH_3	B(OH)_2	H	H
<i>m</i> -BMQBA	CH_3	H	B(OH)_2	H
<i>p</i> -BMQBA	CH_3	H	H	B(OH)_2
BMQ	CH_3	H	H	H

FIGURE 2




FIGURE 3

4/28




FIGURE 4

5/28




FIGURE 5

6/28




FIGURE 6




FIGURE 7




FIGURE 8

Table 1 - Dissociation constants, K_D (μM^3), for the probes with cyanide in water.

Probe	K_D (μM^3)
<i>o</i> -BMOQBA	52.9
<i>m</i> -BMOQBA	84.0
<i>p</i> -BMOQBA	20.8
BMOQ	---
<i>o</i> -BMQBA	16.7
<i>m</i> -BMQBA	16.9
<i>p</i> -BMQBA	15.9
BMQ	---

Table 2 – Multiexponential Intensity decay of BMOQ and o-BMOQBA

[Cyanide] μM	τ_1 (ns)	α_1	τ_2 (ns)	α_2	$\bar{\tau}$ (ns)	$\langle\tau\rangle$ (ns)	χ^2
*o-BMOQBA							
0	26.71	1.0			26.71	26.71	1.33
5	26.33	1.0			26.33	26.33	1.13
10	26.34	1.0			26.34	26.34	1.21
15	26.19	1.0			26.19	26.19	1.30
25	24.78	1.0			24.78	24.78	1.23
35	0.324	0.0160	25.54	0.9840	25.53	25.14	1.35
45	0.326	0.0184	25.10	0.9816	25.09	24.64	1.46
50	0.455	0.0176	25.20	0.9824	25.19	24.76	1.41
*BMOQ							
0	27.30	1.0			27.30	27.30	1.08
5	27.04	1.0			27.04	27.04	1.10
10	26.74	1.0			26.74	26.74	1.12
15	26.53	1.0			26.53	26.53	1.06
20	26.25	1.0			26.25	26.25	1.14
30	25.86	1.0			25.86	25.86	1.17
40	25.37	1.0			25.37	25.37	1.05
50	25.00	1.0			25.00	25.00	1.16

* $\lambda_{\text{ex}} = 372 \text{ nm}$, emission was collected with a 416 nm cut-off filter. BMOQ $K_{\text{SV}} \approx 2 \text{ nM}^{-1}$.

FIGURE 10


Table 3 – Multiexponential Intensity decay of BMQ and o-BMQBA

[Cyanide] μM	τ_1 (ns)	α_1	τ_2 (ns)	α_2	$\bar{\tau}$ (ns)	$\langle\tau\rangle$ (ns)	χ^2
*o-BMQBA							
0	2.18	0.4646	4.74	0.5354	4.01	3.55	1.00
5	2.14	0.4615	4.45	0.5385	3.78	3.38	1.12
10	2.28	0.5704	4.75	0.4296	3.78	3.34	1.04
15	1.86	0.3265	3.64	0.6735	3.29	3.06	0.97
20	1.88	0.3476	3.69	0.6524	3.30	3.06	1.04
30	1.44	0.1762	3.27	0.8238	3.11	2.95	1.21
40	1.92	0.3511	3.59	0.6489	3.21	3.00	0.90
50	1.87	0.3320	3.58	0.6680	3.22	3.01	1.07
*BMQ							
0	2.59	1.0			2.59	2.59	1.07
5	2.58	1.0			2.58	2.58	1.09
10	2.59	1.0			2.59	2.59	1.07
15	2.57	1.0			2.57	2.57	1.02
20	2.57	1.0			2.57	2.57	1.12
30	2.55	1.0			2.55	2.55	1.08
40	2.55	1.0			2.55	2.55	1.14
50	2.55	1.0			2.55	2.55	1.17

* $\lambda_{\text{ex}} = 372 \text{ nm}$, emission was collected with a 416 nm cut-off filter. BMQ $K_{\text{sv}} \approx 0.4 \text{ nM}^{-1}$.

FIGURE 11

12/28



Probe	R^1	R^2	R^3
<i>o</i> -BAQBA	B(OH)_2	H	H
<i>m</i> -BAQBA	H	B(OH)_2	H
<i>p</i> -BAQBA	H	H	B(OH)_2
BAQ	H	H	H

FIGURE 12








FIGURE 13

14/28

A



B



C




FIGURE 14




FIGURE 15




FIGURE 16

Table 4: Multiexponential intensity decay of BAQ and o-BAQBA

[Cyanide] μM	τ_1 (ns)	α_1	τ_2 (ns)	α_2	τ_3 (ns)	α_3	$\bar{\tau}$	$\langle\tau\rangle$	χ^2
BAQ									
0	2.48	1	-	-	-	-	2.48	2.48	1.10
2	2.48	1	-	-	-	-	2.48	2.48	1.02
4	2.49	1	-	-	-	-	2.49	2.49	1.19
6	2.49	1	-	-	-	-	2.49	2.49	1.32
10	2.49	1	-	-	-	-	2.49	2.49	1.18
16	2.49	1	-	-	-	-	2.49	2.49	1.28
20	2.47	1	-	-	-	-	2.47	2.47	0.89
o-BAQBA									
(380 nm)^a									
0	2.04	0.71	3.41	0.29	-	-	2.59	2.44	1.06
2	2.02	0.68	3.367	0.32	-	-	2.61	2.45	0.99
4	1.98	0.67	3.37	0.33	-	-	2.61	2.44	0.94
6	1.92	0.62	3.23	0.38	-	-	2.59	2.42	1.06
8 ^c	1.55	0.41	2.98	0.59	-	-	2.60	2.39	1.53
10 ^c	0.67	0.19	2.64	0.81	-	-	2.53	2.27	2.15
12.5	0.44	0.22	2.60	0.78	-	-	2.50	2.12	2.37
	0.21	0.17	2.07	0.63	3.99	0.20	2.76	2.14	1.08
15	0.38	0.28	2.61	0.72	-	-	2.49	1.98	2.18
	0.21	0.23	1.85	0.44	3.46	0.32	2.71	1.97	1.01
20	0.38	0.30	2.65	0.70	-	-	2.52	1.97	2.47
	0.19	0.24	1.69	0.39	3.36	0.37	2.72	1.95	1.12
(550 nm)^b									
0	1.99	0.63	3.19	0.37	-	-	2.57	2.43	0.99
2	1.93	0.59	3.15	0.41	-	-	2.58	2.43	0.98
4	2.04	0.70	3.39	0.30	-	-	2.60	2.45	1.07
6	1.87	0.51	2.97	0.49	-	-	2.53	2.41	1.10
8	1.86	0.55	3.14	0.45	-	-	2.60	2.44	1.01
10	1.75	0.48	3.10	0.52	-	-	2.63	2.45	1.17
12.5	1.85	0.61	3.48	0.39	-	-	2.74	2.49	1.03
15	1.32	0.31	2.93	0.69	-	-	2.66	2.43	1.25
20	1.19	0.30	2.97	0.70	-	-	2.71	2.44	0.92

^a380 nm long-pass filter.^b550±10 nm interference filter.^cNo notable improvement in fit could be obtained using a 3-exponent function. Similar values were also found for the *meta*- and *para*-BAQBA probes.

18/28




FIGURE 18





FIGURE 19

20/28





FIGURE 20




FIGURE 21




FIGURE 22

23/28




FIGURE 23

24/28




FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28